

Caso para diagnóstico

Antonio J. Rondón Lugo

Cordinador comité de Bioética

Instituto de Biomedicina

Universidad Central de Venezuela

Caracas – Venezuela


www.antoniorondonlugo.com

rondonlugo@yahoo.com

Caso Clínico

Niña de 8 años de edad ,con leishmaniasis tegumentaria americana que la tratamos con Glucantime dos años antes y curó . Tenía frotis positivo para leishmanias y leishmanina de 22 mm . En la histopatología se mostraba un granuloma por agente vivo . Se mantuvo asintomática y apareció posteriormente una lesión redondeada alrededor de la cicatriz ,sin prurito u otras manifestaciones.







Diagnóstico Clínico

- ü 1.Psoriasis
- ü 2.Liquen Plano
- ü 3.Leishmaniasis
- ü 4.Cromomicosis
- ü 5.Tuberculosis Cutánea

Que exámenes practicaría

- n 1.Leishmanina
- n 2.Esporotriquina
- n 3.Examen Micológico : Directo-Cultivo
- n 4.Cultivo para micobacterias
- n 5.Biopsia






Diagnóstico. ?...

Diagnóstico ?

- n 1.Leishmaniasis ?
- n 2.Cromomicosis ?
- n 3.Esporotricosis ?
- n 4.Tuberculosis ?



Diagnóstico definitivo

Leishmaniasis Cutánea
Intermedia
(Recidivante)

Leishmaniasis Tegumentaria



LCL



C57BL/6
IFN- γ



LCD



BALB/c
IL-4



DBA
IFN γ / IL-4

Formas Intermedias



LCM



LCI

Espectro de la Leishmaniasis

- n ESPECTRO CLÍNICO**
- n ESPECTRO HISTOPATOLÓGICO**
- n ESPECTRO INMUNOLÓGICO**

Espectro de la Leishmaniasis


- n PARÁSITOS**
- n CONDICIONES INMUNOLÓGICAS**
- n CONDICIONES AMBIENTALES**
- n NUMERO DE LOCALIZACIÓN DE PICADURAS**
- n CONDICIONES NUTRICIONALES**
- n INFECCIÓN BACTERIANA**

Espectro: Clínico-HP-Inmunológico


Convit J, Pinardi ME.


Cutaneous Leishmaniasis :the clinical and immunopathological Spectrum .Ciba Foundation Symposium
Rondón Lugo AJ., Convit J.


Spectrum of American Cutaneous Leishmanisis". Dermatology
in five Continents Springer Verlag. Berlin 1988; 789-792











Espectro inmunológico



Convit J, Pinardi ME.

Cutaneous Leishmaniasis :the clinical and immunopathological Spectrum .Ciba Foundation Symposium
Rondón Lugo AJ., Convit J.

Spectrum of American Cutaneous Leishmanisis". Dermatology
in five Continents Springer Verlag. Berlin 1988; 789-792

Convit J., Ulrich M., Tapia FJ., Castes M., Rondón Lugo AJ

The Clinical and Immunological Spectrum of American
Cutaneous Leishmanisis". Trans. Roy. Soc. Trop. Medic. 1993. 87: 444-8



Rondón Lugo AJ., Convit J.

“Spectrum of American Cutaneous
Leishmanisis”. Dermatology in five
Continents Springler Verlag. Berlin
1988; 789-792

Halmai O., Arosemena R., Paez E., Weiss E., Rondón Lugo AJ.
“American Cutaneous Leishmaniasis Intermediate Form”. Int. J.
of Dermat. 1993, 32:204-5



*Halmai O., Arosemena R., Paez E.,
Weiss E., Rondón Lugo AJ.*

“American Cutaneous Leishmanisis
Intermediate Form”. Int. J. of
Dermat. 1993, 32:204-5

*Convit J., Ulrich M., Tapia FJ., Castes
M., Rondón Lugo AJ.*

“The Clinical and Immunological Spectrum of American Cutaneous Leishmaniasis”. Trans. Roy. Soc. Trop. Medic. 1993. 87: 444-8

Intermediate or chronic cutaneous leishmaniasis: leukocyte immunophenotypes and cytokine characterisation of the lesion

Diaz NL, Zerpa O, Ponce LV, Convit J, Rondon AJ, Tapia FJ.
Intermediate or chronic cutaneous leishmaniasis: leukocyte immunophenotypes and cytokine characterisation of the lesion.
Exp Dermatol 2002; 11: 34–41. © Munksgaard, 2002

Abstract: The American cutaneous forms of leishmaniasis include immune-responder individuals with localised cutaneous leishmaniasis (LCL) and non-responder individuals with diffuse cutaneous leishmaniasis (DCL). Patients with intermediate or chronic cutaneous leishmaniasis (ICL) have increased morbidity due to the length of their illness, atypical forms and areas of compromise. In the present study, we evaluated the expression of the leukocyte antigens (CD4, CD8, CLA:

**N. L. Diaz, O. Zerpa, L. V. Ponce,
J. Convit, A. J. Rondon and
E. J. Tapia**

Instituto de Biomedicina, Universidad Central de Venezuela, Apartado 4043, Caracas 1010A, Venezuela.

Leishmaniasis Intermedia



LI verrugosa







Leishmaniasis Intermedia



Reprinted from
TRANSACTIONS OF THE ROYAL SOCIETY OF TROPICAL MEDICINE AND HYGIENE.
Vol. 66. No. 4. pp. 603-610, 1972.

DIFFUSE CUTANEOUS LEISHMANIASIS: A DISEASE DUE TO AN IMMUNOLOGICAL DEFECT OF THE HOST

J. CONVIT, M. E. PINARDI AND A. J. RONDÓN


*Division de Dermatología Sanitaria (M. S. A. S.) and Department of Dermatology, Escuela
de Medicina Vargas, Universidad Central de Venezuela, Caracas*



1948. Convit, Lapenta

Grupo Internacional ATD

Invitación



www.buenosalrededor2009.com

•
•
•

Iglesia Clarines



www.antoniorondonlugo.com